

1 2 **Chapter 9 – Geologic Time**3 

- The Importance of a Time Scale
 - Interpreting Earth's history is an important goal of the science of geology
 - Rocks record geologic and evolutionary changes throughout Earth's history
 - Without a time perspective, these events have very little meaning

4 

- Numerical and Relative Dates
 - Numerical dates specify the number of years that have passed since an event occurred
 - Example: The limestone is 250 million years old
 - Prior to the discovery of radioactivity, geologists had no reliable method for numerical dating
 - Relative dates place rocks in a *sequence of formation*
 - Example: The Hermit Shale is older than the Coconino Sandstone
 - Uses a few basic principles, still accurate today

5 

- Principle of Superposition
 - In an undeformed sequence of sedimentary rocks, each bed is older than the one above and younger than the one below
 - This principle also applies to surface features like lava flows and beds of ash
 - Developed by Steno in 1669

6 **Superposition**7 

- Principle of Original Horizontality
 - Layers of sediment are generally deposited in a horizontal position
 - Rock layers that are flat have not been disturbed
 - Rock layers that are deformed, must have been *deformed after deposition*

8 **Original Horizontality**9 

- Principle of Lateral Continuity
 - Beds originate as continuous layers that extend in all directions until they eventually thin out or grade into a different sediment type
 - When a river carves a canyon, we can assume that similar strata on either side were once connected across the span of the canyon

10 **Lateral Continuity**11 **Lateral Continuity**12 

- Principle of Cross-Cutting Relationships
 - Younger features cut across older features
 - Features that cut across rocks (*faults*, intrusions) must have formed after the rocks they cut through

13 **Cross-Cutting**14 **Cross-Cutting**15 

- Principle of Inclusions
 - *Inclusions* are fragments of one rock unit that are enclosed within another rock unit
 - The rock containing the inclusion is younger
 - When magma intrudes a rock mass, blocks of that rock may become dislodged and incorporated into the magma
 - These inclusions are called *xenoliths*
 -

16  **Inclusions**17 

- Unconformities
 - Layers of rock that have been deposited without interruption are called conformable layers
 - An unconformity is a break in the rock record produced by nondeposition and erosion of rock units
 - Uplift and erosion is followed by subsidence and renewed deposition
 - Three basic types: angular unconformity, nonconformity, disconformity

18  **Unconformity**19  **Siccar Point, Scotland**20 

- Unconformities
 - Angular unconformity
 - Tilted rocks are overlain by flat-lying rocks
 - Disconformity
 - Sedimentary strata on either side of the unconformity are parallel
 - Nonconformity
 - Sedimentary strata overlay metamorphic or igneous rocks
 - All three types of unconformities can be seen in the Grand Canyon

21  **Unconformity**22  **Unconformity**23  **Unconformity**24  **Fossils: Evidence of Past Life**

- Fossils are traces or remains of prehistoric life preserved in rock
- Paleontology is the study of fossils
- Knowing the nature of life that existed at a particular time helps researchers understand past environmental conditions
 - Fossils play a key role in correlating rocks of similar ages from different places on Earth











25  **Fossils: Evidence of Past Life**

- Types of Fossils
 - Permineralization
 - Mineral-rich groundwater flows through porous tissue (e.g., bone or wood) and precipitates minerals
 - *Petrified* literally means “turned to stone”
 - Molds and Casts
 - A *mold* is created when a shell is buried in sediment and then dissolved by underground water
 - A *cast* is created when the hollow spaces of a mold are filled with mineral matter

26  **Fossils: Evidence of Past Life**

- Types of Fossils
 - Carbonization and Impressions
 - Carbonization occurs when an organism is buried and compressed, which squeezes out gases and liquids leaving a thin film of carbon behind
 - Effective at preserving leaves and delicate animals
 - Impressions remain in the rock when the carbon film is lost
 - Amber
 - Amber is the hardened resin of ancient trees
 - Effective at preserving insects
 - Trace Fossils
 - Indirect evidence of prehistoric life
 - Includes tracks, burrows, coprolites, and gastroliths

27  **Types of Fossils**

- 28  **Fossils: Evidence of Past Life**
- Conditions favoring preservation
 - Most organisms are not preserved, two special conditions are necessary for most fossil preservation:
 - Rapid burial and
 - The possession of hard parts
- 29  **Correlation of Rock Layers**
- Correlation involves matching of rocks of similar ages from different regions
 - Correlation provides a more comprehensive view of the rock record
 - Often accomplished by noting the position of the bed in a sequence of strata
 - Involves matching of rocks of similar ages from different regions
 - To correlate over larger areas, fossils are used for correlation
- 30  **Correlation**
- 31  **Correlation of Rock Layers**
- Fossils and Correlation
 - Principle of Faunal Succession
 - Used by William Smith, British canal builder
 - The principle of fossil succession states that fossils are arranged according to their age
 - Example: Age of Trilobites, Age of Fishes, Age of Reptiles, Age of Mammals
 - Index Fossils and Fossil Assemblages
 - Index fossils are widespread geographically and limited to a short period of geologic time
 - Fossil assemblages can be used to identify a rock bed that does not contain an index fossil
- 32  **Index Fossils**
- 33  **Fossil Assemblage**
- 34  **Correlation of Rock Layers**
- Environmental Indicators
 - Fossils can be used to infer information about past environments
 - Shells of organisms can be used to infer positions of ancient shorelines and seawater temperatures
 - Corals can be used to indicate former temperature of the water
- 35  **Dating with Radioactivity**
- Reviewing Basic Atomic Structure
 - The nucleus contains *protons* and *neutrons*
 - Protons are positively charged particles with mass
 - Neutrons are a combination of a proton and electron and have a neutral charge
 - *Electrons* are negatively charged particles that orbit the nucleus
 - The *atomic number* is the number of protons in the nucleus
- 36  **Dating with Radioactivity**
- Reviewing Basic Atomic Structure
 - The mass number is the number of protons and neutrons in a nucleus
 - Isotopes have
 - Same number of protons
 - Different numbers of neutrons
 - Different atomic mass
- 37  **Dating with Radioactivity**
- Radioactivity is the spontaneous decay in the structure of an atom's nucleus
 - Types of radioactive decay
 - Alpha emission—an *alpha particle* (two protons and two neutrons) are ejected from the atom
 - Mass number is reduced by 4, and the atomic number is lowered by 2

- Beta emission—a *beta particle* (an electron) is ejected from the atom
 - A neutron is composed of an electron and a proton. When the electron is ejected, the mass number remains unchanged and the atomic number is increased by 1
- Electron capture—an electron is captured in the nucleus
 - The electron combines with the proton and changes into a neutron. The mass number remains unchanged and the atomic number is decreased by 1

38 **Dating with Radioactivity**

- Radioactivity
 - Unstable radioactive isotope is called the *parent*
 - Isotopes resulting from the decay of a parent are termed the *daughter products*
 - The ratio between parent and daughter isotopes in a rock is used to determine its numerical age

39 **Common Types of Radioactive Decay**

40 **Radioactive Decay**

41 **Dating with Radioactivity**

- Radioactivity
 - Radiometric dating
 - Uses the decay of isotopes in rocks to calculate the age of that rock
- Half-Life
 - A half-life is the amount of time required for half of the radioactive isotope to decay
 - Radioactive parent isotopes decay to stable daughter isotopes
 - When the ratio of parent to daughter is 1:1, one half-life has passed

42 **Radioactive Decay Curve**

43 **Dating with Radioactivity**

- Using Various Isotopes
 - With each passing half-life, 50 percent of the remaining parent decays to daughter atoms
 - As the parent atoms decrease, the daughter atoms increase
 - Several naturally occurring radioactive isotopes are useful for dating rocks
 - Potassium-argon: commonly used example
 - Has a half-life of 1.3 billion years
 - Can date rocks as young as 100,000 years
 - Potassium-40 (^{40}K) decays to argon-40 (^{40}Ar) and calcium-40 (^{40}Ca)
 - ^{40}Ar is a gas and only present in rocks as the daughter product of the decay of ^{40}K

44 **Isotopes Frequently Used in Radiometric Dating**

45 **Dating with Radioactivity**

- A Complex Process
 - Determining the quantities of parent and daughter isotopes must be precise
 - Some radioactive materials do not decay directly into stable daughter isotopes
 - Uranium-238 has 14 steps to ultimately decay to the stable daughter lead-206

46 **Dating with Radioactivity**


- Sources of Error
 - The system must be closed
 - No external addition or loss of parent or daughter isotopes
 - Fresh, unweathered rocks are ideal to use for radiometric dating
- Earth's Oldest Rocks
 - Oldest rocks are found on the continent
 - All continents have rocks exceeding 3.5 billion years
 - Confirms the idea that geologic time is immense

47 **Dating with Radioactivity**

- Dating with Carbon-14
 - Radiocarbon dating uses the radioactive isotope carbon-14 to date geologically recent events
 - The half-life of carbon-14 is 5730 years
 - Can be used to date events from the historic past to events as old as 70,000 years
 - Carbon-14 is produced in the upper atmosphere from cosmic-ray bombardment
 - Carbon-14 is incorporated into carbon dioxide and absorbed by plants through photosynthesis
 - *Carbon-14 is only useful in dating organic matter*
 - » All organisms contain a small amount of carbon-14

48  **Carbon-14**49  **The Geologic Time Scale**


- The geologic time scale encompasses all of Earth's history
 - Subdivides geologic history into units with meaningful time frames
 - Originally created using relative dates
 - Numerical dates applied to it in the twentieth century

50  **Geologic Time Scale**51  **The Geologic Time Scale**

- Structure of the Time Scale
 - An eon represents the greatest expanse of time
 - The Phanerozoic eon (“visible life”) is the most recent eon, which began about 542 million years ago.
 - Eons are divided into eras
 - The Phanerozoic eon is divided into three eras
 - Paleozoic era (“ancient life”)
 - Mesozoic era (“middle life”)
 - Cenozoic era (“recent life”)

52  **The Geologic Time Scale**


- Structure of the Time Scale (continued)
 - Each Phanerozoic era is divided into periods
 - The Paleozoic era has seven periods
 - The Mesozoic and Cenozoic eras each have three periods
 - Each period is divided into epochs
 - Except for the seven recent epochs in the Cenozoic, most epochs are termed early, middle, and late

53  **The Geologic Time Scale**

- Precambrian Time
 - Most detail in the geologic time scale is in the Phanerozoic eon
 - The 4 billion years (88% of Earth's history) prior to the Cambrian period are divided into two eons and often collectively referred to as the Precambrian
 - Proterozoic – “Before Life”
 - Archean – “Ancient”
 - Less is known about Earth further back in geologic time

54  **The Geologic Time Scale**

- Precambrian Time (continued)
 - During the Precambrian, simple life-forms that lacked a hard part (algae, bacteria, worms, fungi) dominated
 - First abundant fossil evidence does not appear until the beginning of the Cambrian period
 - Many Precambrian rocks are highly deformed metamorphic rocks—destroying any evidence of past environments

55  **The Geologic Time Scale**

- Terminology and the Geologic Time Scale
 - *Precambrian* is an informal name for the eons before the Phanerozoic
 - *Hadean* refers to the earliest interval of Earth's history—BEFORE the oldest known rocks
 - Geologic timescale is continuously updated
 - *Anthropocene*—referring to human history—is suggested to identify the time since the 1800s that the Earth has seen human-caused global environmental change

56  **Determining Numerical Dates for Sedimentary Strata**

- Sedimentary rocks can rarely be dated directly by radiometric means
 - Geologists must rely on igneous rocks in the strata
 - Radiometric dating determines the age of the igneous rocks
 - Relative dating techniques assign date ranges to sedimentary rocks
 - This is referred to as “bracketing” various episodes in Earth's history
 - Shows the necessity of combining laboratory dating methods with relative dating principles

57  **Dating Sedimentary Strata**

58 

End of Chapter 9